

**Glacier response to North Atlantic climate variability during the Holocene**

N. L. Balasacio et al.

# Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio<sup>1</sup>, W. J. D'Andrea<sup>1</sup>, and R. S. Bradley<sup>2</sup>

<sup>1</sup>Lamont-Doherty Earth Observatory of Columbia University, Palisades, NY 10964, USA

<sup>2</sup>Department of Geosciences, University of Massachusetts, Amherst, MA 01003, USA

Received: 16 April 2015 – Accepted: 28 April 2015 – Published: 27 May 2015

Correspondence to: N. L. Balasacio (balasacio@ldeo.columbia.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Small glaciers and ice caps respond rapidly to climate variations and records of their past extent provide information on the natural envelope of past climate variability. Millennial-scale trends in Holocene glacier size are well documented and correspond with changes in Northern Hemisphere summer insolation. However, there is only sparse and fragmentary evidence for higher frequency variations in glacier size because in many Northern Hemisphere regions glacier advances of the past few hundred years were the most extensive and destroyed the geomorphic evidence of ice growth and retreat during the past several thousand years. Thus, most glacier records have been of limited use for investigating centennial scale climate forcing and feedback mechanisms. Here we report a continuous record of glacier activity for the last 9.5 ka from southeast Greenland, derived from high-resolution measurements on a proglacial lake sediment sequence. Physical and geochemical parameters show that the glaciers responded to previously documented Northern Hemisphere climatic excursions, including the “8.2 ka” cooling event, the Holocene Thermal Maximum, Neoglacial cooling, and 20th Century warming. In addition, the sediments indicate centennial-scale oscillations in glacier size during the late Holocene. Beginning at 4.1 ka, a series of abrupt glacier advances occurred, each lasting ~ 100 years and followed by a period of retreat, that were superimposed on a gradual trend toward larger glacier size. Thus, while declining summer insolation caused long-term cooling and glacier expansions during the late Holocene, climate system dynamics resulted in repeated episodes of glacier expansion and retreat on multi-decadal to centennial timescales. These episodes coincided with ice rafting events in the North Atlantic Ocean and periods of regional ice cap expansion, which confirms their regional significance and indicates that considerable glacier activity on these timescales is a normal feature of the cryosphere. The data provide a longer-term perspective on the rate of 20th century glacier retreat and indicate that recent anthropogenic-driven warming has already impacted the regional cryosphere in a manner outside the natural range of Holocene variability.

## Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



## 1 Introduction

Glaciers and ice caps represent a small but important portion of the cryosphere (~ 785 000 km<sup>2</sup>; Dyurgerov and Meier, 2005). Their mass wasting during the 20–21st century is responsible for 60 % of the sea-level rise unattributable to ocean warming (Meier, 2007) and they continue to retreat at an exceptional rate (Zemp et al., 2012). Moreover, because small glaciers and ice caps respond rapidly to climate changes and there is a strong relationship between glacier mass balance and summer temperature (Oerlemans, 2005), past glacier extent can inform us about past climate variability.

Holocene glacier activity in the Arctic is reasonably well documented at millennial timescales (Miller et al., 2010). Northern Hemisphere glaciers receded in the early Holocene and were smaller than present during the mid-Holocene. Centennial-scale variations, however, are not well constrained because there are few high-resolution and continuous records, and because in many regions the most extensive glacier advances since the early Holocene took place within the past few hundred years, destroying evidence of intervening glacier positions. This is generally the case in Greenland, where historical (AD 1200–1940) advances of local glaciers were generally the most extensive since at least the early Holocene (Kelly and Lowell, 2009).

Evidence from the Greenland Ice Sheet (Kobashi et al., 2011), marine sediments (Bond et al., 1997; Thornalley et al., 2009; Moffa-Sánchez et al., 2014; Jiang et al., 2015), and terrestrial archives (D’Andrea et al., 2011; Larsen et al., 2012; Olsen et al., 2012) indicate that abrupt changes in atmospheric circulation and ocean dynamics, including abrupt cooling events, have punctuated the Holocene. These episodes have alternately been attributed to solar variability, freshwater forcing, volcanic activity, and/or changes in Atlantic Meridional Overturning Circulation (Wanner et al., 2011). How sensitive were glaciers to these abrupt episodes, and did glaciers throughout the North Atlantic respond uniformly? During the period from AD 1250–1900, often referred to as the “Little Ice Age,” well-resolved records from the North Atlantic region suggest coherence in ice cap activity that was potentially driven by volcanic activity coupled

CPD

11, 2009–2036, 2015

### Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with sea-ice/ocean feedbacks (Miller et al., 2012). However, prior to the last 1000 years there are sparse data for use in investigating the synchrony of glacier response to climate variability in the North Atlantic region.

Lakes that receive meltwater from temperate glaciers can be used to develop continuous records of glacier activity. Bedrock erosion at the base of glaciers provides sediment supply for meltwater transport to proglacial lakes. In catchments where other sources of sediment are limited, such as from mass wasting or the release of stored sediment, there is a strong relationship between sediment properties and glacier size (Nesje et al., 2000; Dahl et al., 2003; Jansson et al., 2005), which also follows from the assumption that large glaciers produce more minerogenic material and meltwater than small glaciers. Measurements of physical and geochemical properties of proglacial lake sediments can therefore be used to reconstruct records of past glacier size. Here we report a continuous 9.5 ka record of glacier activity on Kulusuk Island, southeast Greenland developed using sediment cores recovered from Kulusuk Lake (65.56° N, 37.11° W; 202 m) (Fig. 1).

## 2 Study site

Kulusuk Lake (0.8 km<sup>2</sup>, 69 m maximum depth) is located below a cirque with two small glaciers and is within a low arctic maritime region (MAT -1 °C, MAP 900 mm). Characteristic erosional features indicate that local glaciers have temperate thermal structures (Humlum and Christiansen, 2008). Distinct moraines defined by sharp crests are located in front of both glaciers and the recently glaciated area accounts for ~ 50 % of the catchment, which is composed of Archaen gneisses (Bridgwater, 1976) (Fig. 1). Kulusuk Lake is ideally situated to capture and preserve a clear sedimentary record of glacier activity because: (i) it only receives runoff from a very small catchment (catchment : lake area ratio of ~ 2 : 1), minimizing the potential for long-term storage of sediments prior to deposition and limiting sediment input from non-glacial processes, (ii) the proximity of the glaciers to the lake results in minimal sediment transport distance,

CPD

11, 2009–2036, 2015

## Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and (iii) the small size of the glaciers makes them sensitive to minor climate variations (Fig. 1). Therefore, bedrock erosion by the glaciers provides the primary source of minerogenic sediment to the lake and changes in glacier size should clearly be reflected in sediment properties.

## 3 Methods

### 3.1 Sediment core collection and analysis

Sediment cores were recovered from Kulusuk Lake in April 2010 when the lake was ice covered. Bathymetric measurements were made manually through holes drilled in the ice and sediment cores were collected using Uwitec gravity and percussion coring devices from the deepest location, which has a water depth of 69 m. A composite 3.5 m record was compiled by matching the physical stratigraphy and scanning XRF profiles from a 26 cm gravity core (Kul-10D-B) and multiple overlapping percussion cores (Kul-10G-A1, -B1, -A2).

The magnetic susceptibility of the cores was measured every 0.5 cm using a Bartington MS2E sensor. The organic-matter content of the cores was measured by loss-on-ignition (LOI) on contiguous 1 cm<sup>3</sup> samples taken at 1 cm intervals. Organic-matter content was calculated as the difference between the weight of dried 1 cm<sup>3</sup> samples and their weight after heating for 4 h at 550 °C (Dean, 1974). Bulk density measurements (g cm<sup>-3</sup>) and the calculated sedimentation rates (cm year<sup>-1</sup>) were used to determine mass accumulation rates (MAR; g cm<sup>-2</sup> year<sup>-1</sup>). Grain size measurements were made at 10 cm increments. Samples were pre-treated with a 30% hydrogen peroxide solution to digest organic material and analyzed using a Beckman Coulter LS200 particle-size analyzer.

CPD

11, 2009–2036, 2015

## Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 3.2 Chronology

An age-depth model was established based on  $^{210}\text{Pb}$  analysis of the upper sediments and AMS radiocarbon dates on macrofossils. The  $^{210}\text{Pb}$  activity of samples taken every 1 cm from the upper 10 cm of the record was measured by Flett Research Ltd. (Winnipeg, Canada) and ages were modeled from these data using a constant rate of supply model. AMS radiocarbon measurements were made on plant/wood fragments and *Daphnia ephippia* that were wet sieved from core samples. All radiocarbon ages were calibrated to calendar years using CALIB v. 6.0 (Stuiver and Reimer, 1993) with the IntCal09 calibration dataset (Reimer et al., 2009). Ages are presented in calendar years prior to AD 1950 (BP) unless otherwise indicated.

## 3.3 Scanning X-ray fluorescence

To characterize minerogenic changes at higher resolution and with greater sensitivity, an Itrax<sup>TM</sup> XRF core scanner was used to produce profiles of relative elemental compositions (Croudace et al., 2006). The Itrax<sup>TM</sup> continuously scans the surface of sediment cores at sub-mm resolution with a micro X-ray beam (20 mm × 100 μm) and the relative concentrations of a range of elements are determined based the detection of dispersive energy spectra. Dispersive energy spectra are acquired across each measured interval and peak area integrals are calculated for each element. Peak area integrals are related to elemental concentrations within the sediment, but can also be influenced by characteristics of the sedimentary matrix and therefore only indicate relative changes in elemental composition (Croudace et al., 2006; Rothwell et al., 2006). Our analysis focused on the elements: K, Ca, Ti, Mn, Fe, Zn, Rb, Sr, which have detection limits that range from 150 to 5 ppm (Croudace et al., 2006). All of the cores were scanned at 200 μm intervals with an exposure time of 10 s, voltage of 30 kV, and current of 55 nA.

CPD

11, 2009–2036, 2015

## Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





increase and then display more minor fluctuations across the upper 1.2 m. These intervals are clearly defined by MARs, which incorporate sediment density measurements that range from  $\sim 0.8\text{--}1.8\text{ g cm}^{-3}$ , but are primarily controlled by the large sedimentation rate changes (Fig. 3).

## 4.2 Scanning XRF data analysis

Elemental scans acquired by scanning XRF show a similar response to magnetic susceptibility with higher values across coarser, clastic intervals. However, the XRF data have a greater sensitivity to minerogenic changes and were measured at higher resolution (0.2 mm) (Fig. 3). We focused our analysis on the elements: K, Ca, Ti, Mn, Fe, Zn, Rb, and Sr, which are common in siliciclastic sediments. Changes in the concentrations of these elements reflect changes in the contribution of minerogenic material eroded from catchment bedrock and delivered to the lake. Statistical analysis of the scanning XRF data indicate that all of the elements are highly correlated and that there is a strong primary trend in the data. Correlation coefficients show the strong significant relationships among the majority of the elements (Table 2). Rather than relying on a single element (e.g., Ti), we used principal component analysis (PCA) to define the leading mode of variability (PC1) among the elemental data. PCA allows for a multi-dimensional examination of the dataset in order to identify the primary signal(s). PCA results indicate that there is one strong primary trend in the elemental data with the first eigenvector (PC1) accounting for 76% of the total variance. The factor loadings reveal the high correlations between individual element profiles and PC1 (Table 2). The trends in PC1 are similar to those in the lower resolution magnetic susceptibility and organic matter content records, justifying use of PC1 data to infer past minerogenic changes (Fig. 3). The choice to use PC1 rather than a single representative element (e.g. Ti) to represent changes in sedimentary minerogenic content has no impact on any of our conclusions.

## Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## 5 Discussion

### 5.1 Sedimentation in proglacial lakes

Sedimentation in proglacial lakes can be the result of a complex set of physical processes associated with the erosion, storage, and transport of sediment within glacial and proglacial systems (Dahl et al., 2003; Jansson et al., 2005). It is important to consider these complicating factors when selecting sites for glacier reconstructions and when interpreting sedimentary records. Glaciers fundamentally impact the amount and character of minerogenic sediment in proglacial lakes. Glacier size, erosive ability, and meltwater production directly influence the amount of minerogenic sediment delivered to a proglacial lake. However, sedimentation processes in a proglacial lake can also be impacted by mass wasting processes in paraglacial environments (particularly in landscapes with steep unstable slopes), and by the delayed release of sediment stored along the transport pathway between the glacier and the lake (for example, sediment stored in extensive meltwater stream channels). Relative to minerogenic material, organic sedimentation is typically a minor component in proglacial lakes and is related to the input of organic matter from autochthonous and allochthonous primary productivity, and the preservation thereof. In proglacial environments in the Arctic, low temperatures restrict vegetation and soil cover, and minerogenic sediment input to lakes from glacial meltwater results in turbidity that impedes autochthonous productivity.

Techniques for analyzing sediment from proglacial lakes therefore focus on investigating changes in the character of minerogenic sediment. The minerogenic content of lake sediments, used as a proxy for glacier size, is commonly evaluated by measuring the magnetic susceptibility and organic matter content of the sediments. Magnetic susceptibility reflects the amount of magnetic minerals eroded and input to a lake, and organic matter content is a function of dilution by minerogenic input, changes in primary productivity, and preservation.

At Kulusuk Lake, processes that can complicate the mechanistic link between minerogenic input and glacier size are fundamentally limited. Input of sediment from

## Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



---

**Glacier response to North Atlantic climate variability during the Holocene**

---

N. L. Balasacio et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

non-glacial processes is restricted due to the small catchment and small catchment to lake-area ratio ( $\sim 2 : 1$ ), and the proximity of the glaciers to the lake. These factors also limit the potential for sediment storage between the glaciers and the lake. Furthermore, the landscape surrounding the lake is composed of shallow, low-elevation slopes that minimize the likelihood of mass wasting events (which, if present, would have been easily identified by rapid changes in sediment accumulation, or age reversals, based on  $^{14}\text{C}$  dates). Therefore, at Kulusuk Lake, it is reasonable to interpret changes in minerogenic content as a function of glacier size.

Variations in magnetic susceptibility, organic matter content, and scanning XRF data (PC1) represent changes in the relative amount and grain size of minerogenic sediment delivered to Kulusuk over the last 9.5 ka (Fig. 3). Magnetic susceptibility and PC1 are directly related to, and organic content is inversely proportional to, minerogenic content. The highest magnetic susceptibility and PC1 values correspond to intervals with coarser grain size. We therefore interpret the sedimentological system in Kulusuk Lake as follows: during periods of increased glacier size, more coarse minerogenic sediment was eroded from the bedrock and delivered to the lake by meltwater; during periods of smaller glacier size, less minerogenic sediment was deposited and a greater relative proportion of organic matter content accumulated.

## 5.2 Holocene glacier fluctuations in southeast Greenland

Dramatic changes in minerogenic input to Kulusuk Lake over the last 9.5 ka reveal that the size of the Kulusuk glaciers has varied significantly throughout the Holocene (Fig. 3). Beginning ca. 8.7 ka, increasing organic matter content and decreasing minerogenic content, inferred from magnetic susceptibility and XRF data, indicates significant retreat of the Kulusuk glaciers, corresponding closely in time with the deglaciation of a nearby inland catchment ca. 8.4 ka (Balasacio et al., 2013), following deglaciation of local coastal areas ca. 11.1–9.5 ka (Long et al., 2008; Roberts et al., 2008). A brief interval of increased minerogenic input shows that this early Holocene retreat was interrupted by an episode of advance at 8.5 ka, coeval with reductions in sea sur-

## Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

face temperatures and bottom water circulation in the subpolar North Atlantic, as seen in high-resolution marine records (Ellison et al., 2006; Kissel et al., 2013). Another abrupt episode of minerogenic input ca. 8.2 ka signifies another glacier advance and occurred contemporaneously with the largest abrupt Holocene climate cooling event inferred from Greenland ice core records (Thomas et al., 2007). The temporal resolution and age control of this section of our record cannot provide new constraints on the exact timing of these events, however it clearly demonstrates the sensitivity of the Kulusuk glaciers to rapid, regional climate events.

Between 7.8–4.1 ka, the Kulusuk glaciers were at their minimum Holocene extent, inferred from low minerogenic content, low MAR, and high organic matter content in the lake sediments (Fig. 3). We interpret this as an interval with little to no glacier ice in the catchment, primarily based the XRF and magnetic susceptibility data, which are lowest and show reduced variability at this time, relative to the last 4.1 ka. This interval is also marked by extremely high organic matter content that is greater than 12 % (with a maximum of 19 %), suggesting that this period was accompanied by an increase in primary productivity due to a lack of input of glacial flour. Magnetic susceptibility remains close to zero throughout the mid-Holocene section of the core and appears insensitive to the minor minerogenic changes inferred from the XRF data. There are two excursions in the PC1 record during this interval (ca. 7.2 and 6.2 ka), which we interpret as sediment influxes from paraglacial activity rather than as glacier advances because they are short-lived and do not match the amplitude of variation observed elsewhere. Thus, this record provides well-dated constraints on the Holocene Thermal Maximum in this area and refines previous estimates for its onset and termination (Kaufman et al., 2004).

At 4.1 ka, a sharp increase in XRF- and MS-inferred minerogenic content and decrease in organic matter content, indicate the glaciers once again grew large enough to contribute minerogenic material to the lake. The regrowth of the Kulusuk glaciers represents the lowering of the regional snowline, and the precise timing could be considered unique to this catchment. However, the timing is contemporaneous with hydro-

## Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



logic changes at nearby Flower Valley Lake, likely related to an increase in the duration of lake ice cover (Balasacio et al., 2013). We propose that this represents significant cooling and the onset of the regional Neoglacial period. The oscillatory and stepwise increase in minerogenic input (decrease in organic matter content) after 4.1 ka suggests that rather than advancing steadily toward their historical extent, the Kulusuk glaciers episodically advanced and retreated at centennial timescales until ca. 1.3 ka. After ca. 1.3 ka they stabilized and advanced only gradually until their rapid 20th century retreat (Fig. 3). Importantly, the major sedimentological transitions in the record are all located near radiocarbon dates, thereby maximizing the certainty of their timing and the calculations of sediment accumulation rates. The timing of glacier size variations between radiocarbon-dated intervals since 4.1 ka are interpolated, and we estimate the accuracy to be better than  $\pm 100$  years, the average 2- $\sigma$  uncertainty of the ages.

### 5.3 Evidence for synchronous regional glacier response during the late Holocene

The Kulusuk glacier reconstruction documents centennial scale episodes of glacier advance during the Neoglacial (4.1 to 1.3 ka) coeval with other records of glacier growth in the North Atlantic region. After 4.1 ka, six major advances of the Kulusuk glaciers occurred (4.1, 3.9, 3.2, 2.8, 2.1, and 1.3 ka) and each successive advance resulted in greater glacier extent (Fig. 4). The progressive increase in glacier size is consistent with declining NH summer insolation. However, each episode of glacier advance was followed by a period of retreat (or at least stabilization), possibly suggesting that the glaciers repeatedly grew out of equilibrium with external insolation forcing and then retreated back toward an equilibrium state. The episodic advances of the Kulusuk glaciers during the past 4.1 ka are nearly identical in timing to the cooling episodes in the North Atlantic Ocean inferred from ice-rafted hematite-stained grains identified in marine sediment cores (Bond et al., 1997, 2001) (Fig. 4). Cooling events at these times have also been documented on the East Greenland and Icelandic Shelves and attributed to increased strength of the East Greenland Current (Giraudeau et al., 2000;

---

**Glacier response to North Atlantic climate variability during the Holocene**

---

N. L. Balasacio et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Jennings et al., 2002; Ran et al., 2008). Moreover, the Langjökull ice cap in Iceland advanced along with the Kulusuk glaciers and the North Atlantic IRD events (Larsen et al., 2012) (Fig. 4), and advances of the Bregne ice cap in east Greenland at ca. 2.6 and 1.9 ka (Levy et al., 2014), within chronological uncertainty of the Kulusuk glacier advances ca. 2.8 and 2.1 ka, have also been documented. We propose that continuous records of glacier activity around the North Atlantic during the Neoglacial are beginning to reveal synchronous glacier response to abrupt episodes of climate change.

Although the amplitude of variability in the proxy measurements during the past 1.3 ka are lower than earlier in the Holocene, due to the greater size and stability of the Kulusuk glaciers, it is still worthwhile to examine the changes in the sediment properties during this time. The very high sediment accumulation rates during this interval ( $0.8 \text{ mm year}^{-1}$ ), allow sub-annual XRF measurements and, if interpreted in the same manner as periods with smaller glacier size, can afford a detailed examination of changes in glacier size using the XRF PC1 data (Fig. 5). The overall trend reveals a small and very gradual glacier expansion after 1.3 ka followed by 20th century retreat, which resembles the overall trend in Arctic temperatures over the last 2 ka (Kaufman et al., 2009).

Multi-decadal variations in inferred glacier size during the past 1.3 ka also appear synchronous with those of other glaciers in the region (Fig. 5). Kulusuk glaciers increased in size ca. AD 1250–1300 and again ca. AD 1350 and AD 1450, precisely when ice caps on Baffin Island (Miller et al., 2012) and Iceland (Larsen et al., 2011) were expanding (Fig. 4). After AD 1450 Kulusuk glaciers continued to expand, as did Langjökull on Iceland, while evidence from the Baffin ice caps indicates continuous ice cover (Miller et al., 2012). Both Kulusuk and Langjökull glaciers appear to have advanced in two phases, at ca. AD 1450–1630 and ca. AD 1700–1930. Varve thickness and magnetic susceptibility data from a proglacial lake on Baffin Island, if interpreted to reflect glacier size, also reveal two similar distinct glacier advances at these times (Fig. 4) as well as the earlier advance ca. AD 1250–1300 also observed in the Kulusuk record (Thomas and Briner, 2009; Thomas et al., 2010). Taken together, these data are

## Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



consistent with records from around Greenland that suggest the most extensive glacier advances since the early Holocene occurred between AD 1250–1900, and provide evidence for regionally coherent cooling phases during the Little Ice Age (Grove, 2001). We note that this timing contrasts with evidence from east Greenland that suggests LIA advance of the Istorvet Ice Cap approximately 100 years earlier (ca. AD 1150; Lowell et al., 2013), unless the data are reinterpreted as suggested by Miller et al. (2013).

Therefore, we propose that there has been regional coherence in glacier activity not only during the past 1.3 ka, as previously suggested (Miller et al., 2012), but also during the past 4.1 ka, and glacier growth in response episodic climate change has been a common feature of glaciers in the North Atlantic region throughout, at least, the last 4.1 ka.

Cold events are an important feature of centennial-scale climate of the Holocene (Wanner et al., 2011). Cooling events in the North Atlantic region are likely directly associated with changes in Atlantic Meridional Overturning Circulation (AMOC) (Denton and Broecker, 2008). IRD data suggest that periodic circulation changes of the North Atlantic Ocean resulted in an advection of cold, fresh surface water south and east during ice-rafting events throughout the Holocene (Bond et al., 1997). Ocean circulation changes associated with IRD events and sea-surface temperatures have been attributed to solar forcing (Bond et al., 2001; Moffa-Sánchez et al., 2014; Jiang et al., 2015) and some modeling studies have confirmed that AMOC can switch between distinct modes in response to a small external forcing, such as solar variability (Jongma et al., 2007). However, modeling results are inconsistent and it is also possible that cooling events might have simply resulted from internal ocean dynamics (Schulz and Paul, 2002). Regardless of the mechanism, our results demonstrate that glaciers responded quite actively to natural climate variations of the Holocene.

### 5.4 Rates of glacier change during the Holocene

This well-dated, high-resolution record of changes in the size of the Kulusuk glaciers also allows comparison among the rates of past glacier size variations. We present

## Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



relative rates of change inferred from the first derivative of the XRF PC1 data in 105 year binned averages, an interval chosen using the interval with the lowest resolution (Fig. 6). We acknowledge the caveat that they are based on the assumption that the relationship between minerogenic input and glacier size has remained constant.

5 The analysis indicates that the rate of 20th century retreat of the Kulusuk glaciers was greater than during any other century of the past 1.3 ka, including during the Medieval Climate Anomaly. Furthermore, the 20th century retreat rate was two to three times the rate of any other period of retreat during the past 4.1 ka, and almost twice as rapid as the early Holocene retreat that marked the transition into the regional HTM (Fig. 6).  
10 This comparison helps to place the rate of 20th century glacier loss in the context of natural episodes of past glacier activity.

## 6 Conclusions

The Kulusuk Lake sediment record was used to generate a high-resolution record of changes in the size of the Kulusuk glaciers over the last 9.5 ka. The record shows that the glaciers were sensitive to a number of previously documented regional climate fluctuations and extends our understanding of Holocene climate dynamics in this sector of the Arctic. In particular, the record clearly constrains the Holocene Thermal Maximum at this site to between 7.8 and 4.1 ka, when the glaciers likely completely melted away. The regrowth of the Kulusuk glaciers at 4.1 ka corresponds with regional hydrologic changes and reflects the onset of the Neoglacial Period. The last 4.1 ka is marked by a series of abrupt glacier advances as the size of the Kulusuk glaciers increased. These episodes of glacier growth correspond with ice rafting events in the North Atlantic Ocean, as well as regional ice cap expansion, and demonstrate that glaciers in this sector of the Arctic were very active during the late Holocene in response to abrupt cooling events that punctuated millennial-scale insolation-driven cooling. The reconstruction of Kulusuk glacier activity provides a new and refined perspective on  
15  
20  
25

late Holocene cold events, which are important features of centennial-scale climate variability.

*Acknowledgements.* This research was supported by a LDEO Postdoctoral Fellowship to NLB, NSF grant ARC-0851642 to WJD, NOAA grant NA09OAR4600215 and NSF grant ARC-0909354 to RSB. We thank Lucien von Gunten, Sam Davin, and Greg de Wet for assistance with field work.

## References

- Balascio, N. L., D'Andrea, W. J., Bradley, R. S., and Perren, B. B.: Biogeochemical evidence for hydrologic changes during the Holocene in a lake sediment record from southeast Greenland, *Holocene*, 23, 1428–1439, 2013.
- Berger, A. and Loutre, M. F.: Insolation values for the climate of the last 10 million years, *Quaternary Sci. Rev.*, 10, 297–317, 1991.
- Bond, G., Showers, W., Cheseby, M., Lotti, R., Almasi, P., deMenocal, P., Priore, P., Cullen, H., Hajdas, I., and Bonani, G.: A pervasive millennial-scale cycle in North Atlantic Holocene and Glacial climates, *Science*, 278, 1257–1266, 1997.
- Bond, G., Kromer, B., Beer, J., Muscheler, R., Evans, M. N., Showers, W., Hoffmann, S., Lotti-Bond, R., Hajdas, I., and Bonani, G.: Persistent solar influence on North Atlantic climate during the Holocene, *Science*, 294, 2130–2136, 2001.
- Bridgwater, D.: Nagssustoqidian mobile belt in East Greenland, *Geology of Greenland*, 7, 97–103, 1976.
- Croudace, I. W., Rindby, A., and Rothwell, R. G.: ITRAX: description and evaluation of a new multi-function X-ray core scanner, in: *New Techniques in Sediment Core Analysis*, edited by: Rothwell, R. G., Special Publications 267, Geological Society, London, 51–564, 2006.
- D'Andrea, W. J., Huang, Y., Fritz, S. C., and Anderson, N. J.: Abrupt Holocene climate change as an important factor for human migration in West Greenland, *P. Natl. Acad. Sci. USA*, 108, 9765–9769, 2011.
- Dahl, S. O., Bakke, J., Lie, Ø., and Nesje, A.: Reconstruction of former glacier equilibrium-line altitudes based on proglacial sites: an evaluation of approaches and selection of sites, *Quaternary Sci. Rev.*, 22, 275–287, 2003.

## Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Dean, W. E.: Determination of carbonate and organic matter in calcareous sediments and sedimentary rocks by loss on ignition: comparison with other methods, *J. Sediment. Petrol.*, 44, 242–248, 1974.

Denton, G. H. and Broecker, W. S.: Wobbly ocean conveyor circulation during the Holocene, *Quaternary Sci. Rev.*, 27, 1939–1950, 2008.

Dyurgerov, M. and Meier, M. F.: *Glaciers and the Changing Earth System: a 2004 Snapshot*, Occasional Paper 58, Institute of Arctic Alpine Research, University of Colorado, Boulder, CO, 118 pp., 2005.

Ellison, C. R. W., Chapman, M. R., and Hall, I. R.: Surface and deep ocean interactions during the cold climate event 8200 years ago, *Science*, 312, 1929–1932, 2006.

Giraudeau, J., Cremer, M., Manthé, S., Labeyrie, L., and Bond, G.: Coccolith evidence for instabilities in surface circulation south of Iceland during Holocene times, *Earth Planet. Sc. Lett.*, 179, 257–268, 2000.

Grove, J. M.: The initiation of the “Little Ice Age” in regions round the North Atlantic, *Climatic Change*, 48, 53–82, 2001.

Humlum, O. and Christiansen, H. H.: Geomorphology of the Ammassalik Island, SE Greenland, *Geogr. Tidsskr.*, 108, 5–20, 2008.

Jansson, P., Rosqvist, G., and Schneider, T.: Glacier fluctuations, suspended sediment flux and glacio-lacustrine sediments, *Geogr. Ann. A*, 87, 37–50, 2005.

Jennings, A. E., Knudsen, K. L., Hald, M., Hansen, C. V., and Andrews, J. T.: A mid-Holocene shift in Arctic sea-ice variability on the East Greenland Shelf, *Holocene*, 12, 49–58, 2002.

Jongma, J. I., Prange, M., Renssen, H., and Schulz, M.: Amplification of Holocene multicentennial climate forcing by mode transitions in North Atlantic overturning circulation, *Geophys. Res. Lett.*, 34, L15706, doi:10.1029/2007GL030642, 2007.

Kaufman, D. S., Ager, T. A., Anderson, N. J., Anderson, P. M., Andrews, J. T., Bartlein, P. J., Brubaker, L. B., Coats, L. L., Cwynar, L. C., Duvall, M. L., Dyke, A. S., Edwards, M. E., Eisner, W. R., Gajewski, K., Geirsdóttir, A., Hu, F. S., Jennings, A. E., Kaplan, M. R., Kerwin, M. W., Lozhkin, A. V., MacDonald, G. M., Miller, G. H., Mock, C. J., Oswald, W. W., Otto-Bliesner, B. L., Porinchu, D. F., Rühland, K., Smol, J. P., Steig, E. J., and Wolfe, B. B.: Holocene thermal maximum in the western Arctic (0–180° W), *Quaternary Sci. Rev.*, 23, 529–560, 2004.

Kaufman, D. S., Schneider, D. P., McKay, N. P., Ammann, C. M., Bradley, R. S., Briffa, K. R., Miller, G. H., Otto-Bliesner, B. L., Overpeck, J. T., Vinther, B. M., and Arctic Lakes 2k Project

## Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Members: recent warming reverses long-term Arctic cooling, *Science*, 325, 1236–1239, 2009.

Kelly, M. A. and Lowell, T. V.: Fluctuations of local glaciers in Greenland during latest Pleistocene and Holocene time, *Quaternary Sci. Rev.*, 28, 2088–2106, 2009.

5 Kissel, C., Van Toer, A., Laj, C., Cortijo, E., and Michel, E.: Variations in the strength of the North Atlantic bottom water during Holocene, *Earth Planet. Sc. Lett.*, 369–370, 248–259, 2013.

Kobashi, T., Kawamura, K., Severinghaus, J. P., Barnola, J.-M., Nakaegawa, T., Vinther, B. M., Johnsen, S. J., and Box, J. E.: High variability of Greenland surface temperature over the past 4000 years estimated from trapped air in an ice core, *Geophys. Res. Lett.*, 38, L21501, doi:10.1029/2011GL049444, 2011.

10 Larsen, D. J., Miller, G. H., Geirsdóttir, Á., and Thordarson, T.: A 3000-year varved record of glacier activity and climate change from the proglacial lake Hvítárvatn, Iceland, *Quaternary Sci. Rev.*, 30, 2715–2731, 2011.

15 Larsen, D. J., Miller, G. H., Geirsdóttir, Á., and Ólafsdóttir, S.: Non-linear Holocene climate evolution in the North Atlantic: a high-resolution, multi-proxy record of glacier activity and environmental change from Hvítárvatn, central Iceland, *Quaternary Sci. Rev.*, 39, 14–25, 2012.

20 Levy, L. B., Kelly, M. A., Lowell, T. V., Hall, B. L., Hempel, L. A., Honsaker, W. M., Lusas, A. R., Howley, J. A., and Axford, Y. L.: Holocene fluctuations of Bregne ice cap, Scoresby Sund, east Greenland: a proxy for climate along the Greenland Ice Sheet margin, *Quaternary Sci. Rev.*, 92, 357–368, 2014.

Long, A. J., Roberts, D. H., Simpson, M. J. R., Dawson, S., Milne, G. A., and Huybrechts, P.: Late Weichselian relative sea-level changes and ice sheet history in southeast Greenland, *Earth Planet. Sc. Lett.*, 272, 8–18, 2008.

25 Lowell, T. V., Hall, B. L., Kelly, M. A., Bennike, O., Lusas, A. R., Honsaker, W., Smith, C. A., Levy, L. B., Travis, S., and Denton, G. H.: Late Holocene expansion of Istorvet ice cap, Liverpool Land east Greenland, *Quaternary Sci. Rev.*, 63, 128–140, 2013.

Meier, M. F.: Glaciers dominate eustatic sea-level rise in the 21st century, *Science*, 317, 1064–1067, 2007.

30 Miller, G. H., Brigham-Grette, J., Alley, R. B., Anderson, L., Bauch, H. A., Douglas, M. S. V., Edwards, M. E., Elias, S. A., Finney, B. P., Fitzpatrick, J. J., Funder, S. V., Herbert, T. D., Hinzman, L. D., Kaufman, D. S., MacDonald, G. M., Polyak, L., Robock, A., Serreze, M. C.,

---

## Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio et al.

---

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Smol, J. P., Speilhagen, R., White, J. W. C., Wolfe, A. P., and Wolff, E. W.: Temperature and precipitation history of the Arctic, *Quaternary Sci. Rev.*, 29, 1679–1715, 2010.

Miller, G. H., Geirsdóttir, Á., Zhong, Y., Larsen, D. J., Otto-Bliesner, B. L., Holland, M. M., Bailey, D. A., Refsnider, K. A., Lehman, S. J., Southon, J. R., Anderson, C., Björnsson, H., and Thordarson, T.: Abrupt onset of the Little Ice Age triggered by volcanism and sustained by sea-ice/ocean feedbacks, *Geophys. Res. Lett.*, 39, L02708, doi:10.1029/2011GL050168, 2012.

Miller, G. H., Briner, J. P., Refsnider, K. A., Lehman, S. J., Geirsdóttir, Á., Larsen, D. J., Southon, J. R.: Substantial agreement on the timing and magnitude of Late Holocene ice cap expansion between East Greenland and the Eastern Canadian Arctic: a commentary on Lowell et al., 2013, *Quaternary Sci. Rev.*, 77, 239–245, 2013.

Moff-Sánchez, P., Born, A., Hall, I. R., Thornalley, D. J. R., and Barker, S.: Solar forcing of North Atlantic surface temperature and salinity over the past millennium, *Nat. Geosci.*, 7, 275–278, 2014.

Nesje, A., Dahl, S. O., Andersson, C., and Matthews, J. A.: The lacustrine sedimentary sequence in Sygneskardvatnet, western Norway: a continuous, high-resolution record of the Jostedalbreen ice cap during the Holocene, *Quaternary Sci. Rev.*, 19, 1047–1065, 2000.

Oerlemans, J.: Extracting a climate signal from 169 glacier records, *Science*, 308, 675–677, 2005.

Olsen, J., Anderson, N. J., Knudsen, M. F.: Variability of the North Atlantic Oscillation over the past 5,200 years, *Nat. Geosci.*, 5, 808–812, 2012.

Ran, L., Jiang, H., Knudsen, K. L., and Eiríksson, J.: A high-resolution Holocene diatom record on the North Icelandic shelf, *Boreas*, 37, 399–413, 2008.

Reimer, P. J., Baillie, M. G. L., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Bronk Ramsey, C., Buck, C. E., Burr, G. S., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Hajdas, I., Heaton, T. J., Hogg, A. G., Hughen, K. A., Kaiser, K. F., Kromer, B., McCormac, F. G., Manning, S. W., Reimer, R. W., Richards, D. A., Southon, J. R., Talamo, S., Turney, C. S. M., van der Plicht, J., and Weyhenmeyer, C. E.: IntCal09 and Marine09 radiocarbon age calibration curves, 0–50,000 years cal BP, *Radiocarbon*, 51, 1111–1150, 2009.

Roberts, D. H., Long, A. J., Schnabel, C., Freeman, S., and Simpson, M. J. R.: The deglacial history of southeast sector of the Greenland Ice Sheet during the Last Glacial Maximum, *Quaternary Sci. Rev.*, 27, 1505–1516, 2008.

## Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Rothwell, R. G., Hoogakker, B., Thomson, J., Croudace, I. W., Frenz, M.: Turbidite emplacement on the southern Balearic Abyssal Plain (western Mediterranean Sea) during Marine Isotope Stages 1–3: an application of ITRAX XRF scanning of sediment cores to lithostratigraphic analysis, in: *New Techniques in Sediment Core Analysis*, edited by: Rothwell, R. G., Special Publications, Geological Society, London, 267, 79–98, 2006.

Schulz, M. and Paul, A.: Holocene climate variability on centennial-to-millennial time scales: 1. Climate records from the North-Atlantic realm, in: *Climate Development and History of the North Atlantic Realm*, edited by: Wefer, G., Berger, W. H., Behre, K.-E., and Jansen, E., Springer, Berlin, 41–54, 2002.

Stuiver, M. and Reimer, P. J.: Extended  $^{14}\text{C}$  database and revised CALIB radiocarbon calibration program, *Radiocarbon*, 35, 215–230, 1993.

Thomas, E. K. and Briner, J. P.: Climate of the past millennium inferred from varved proglacial lake sediments on northeast Baffin Island, Arctic Canada, *J. Paleolimnol.*, 41, 209–224, 2009.

Thomas, E. K., Szymanski, J., and Briner, J. P.: Holocene alpine glaciation inferred from lacustrine sediments on northeastern Baffin Island, Arctic Canada, *J. Quaternary Sci.*, 25, 146–161, 2010.

Thomas, E. R., Wolff, E. W., Mulvaney, R., Steffensen, J., Johnsen, S. J., Arrowsmith, C., White, J. W. C., Vaughn, B., and Popp, T.: The 8.2 ka event from Greenland ice cores, *Quaternary Sci. Rev.*, 26, 70–81, 2007.

Thornalley, D. J. R., Elderfield, H., and McCave, I. N.: Holocene oscillations in temperature and salinity of the surface subpolar North Atlantic, *Nature*, 457, 711–714, 2009.

Wanner, H., Solomina, O., Grosjean, M., Ritz, S. P., and Jetel, M.: Structure and origin of Holocene cold events, *Quaternary Sci. Rev.*, 30, 3109–3123, 2011.

Zemp, M., Frey, H., Gärtner-Roer, I., Nussbaumer, S. U., Hoelzle, M., Paul, F., and Haerberli, W.: *Fluctuations of Glaciers 2005–2010*, Vol. 10, World Glacier Monitoring Service, Zurich, Switzerland, 336 pp., 2010.

## Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 1.** Geochronologic data for the Kulusuk Lake record.

Composite Depth (cm)	Description	Laboratory ID*	<sup>14</sup> C Age (yrBP)	Calibrated Age Range (1-σ) (2-σ)		Median Age (calyrBP)
0	Core Top	–	–	–	–	–60
1	<sup>210</sup> Pb	–	–	–	–	–53
2	<sup>210</sup> Pb	–	–	–	–	–46
3	<sup>210</sup> Pb	–	–	–	–	–36
4	<sup>210</sup> Pb	–	–	–	–	–25
5	<sup>210</sup> Pb	–	–	–	–	–7
6	<sup>210</sup> Pb	–	–	–	–	24
7	<sup>210</sup> Pb	–	–	–	–	44
8	<sup>210</sup> Pb	–	–	–	–	57
9	<sup>210</sup> Pb	–	–	–	–	83
10	<sup>210</sup> Pb	–	–	–	–	111
34	Daphnia ephippia	OS-96479	335 ± 40	316–459	306–486	393
59.5	Plant/wood	UCI-89386	940 ± 20	798–914	795–919	852
95	Daphnia ephippia	OS-96454	1290 ± 25	1183–1276	1178–1283	1237
132	Plant/wood	UCI-87240	3410 ± 60	3574–3814	3484–3832	3664
138.5	Plant/wood	UCI-87241	3820 ± 60	4095–4378	4008–4415	4224
170.5	Daphnia ephippia	OS-96461	7620 ± 50	8377–8452	8359–8539	8418
214.5	Daphnia ephippia	OS-96746	8510 ± 130	9312–9659	9135–9887	9501

\* UCI – University of California Irvine Keck-CCAMS Facility; OS – National Ocean Sciences AMS Facility.

## Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio et al.

**Table 2.** XRF PC1 factor loadings and correlation matrix for scanning XRF elemental data.

	PC1 Loadings	K	Ca	Ti	Mn	Fe	Zn	Rb	Sr
K	0.950	1							
Ca	0.868	0.891	1						
Ti	0.969	0.938	0.797	1					
Mn	0.815	0.672	0.594	0.783	1				
Fe	0.945	0.876	0.748	0.946	0.814	1			
Zn	0.861	0.759	0.609	0.842	0.727	0.836	1		
Rb	0.831	0.743	0.601	0.791	0.637	0.790	0.727	1	
Sr	0.686	0.689	0.803	0.583	0.412	0.488	0.436	0.455	1

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Figure 1.** Location and catchment setting of Kulusuk Lake. The white dashed line marks the watershed boundary and the red dashed line defines the crest of moraines in front of both glaciers. Cores were collected in the deepest basin (red circle). (Image: Google, NASA.)

Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Figure 2.** Magnetic susceptibility profile, percent sand, and mass accumulation rate (MAR) shown next to the age-depth model for the composite Kulusuk Lake record. The four lithostratigraphic units and the corresponding sedimentation rates are shown. A dash line marks the period below the last radiocarbon age (9.5 cal ka BP) where rates of sedimentation are extrapolated.

**Glacier response to North Atlantic climate variability during the Holocene**

N. L. Balasacio et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion







**Figure 3.** Kulusuk Lake record. **(a)** First principal component of the scanning XRF data (PC1). **(b)** Magnetic susceptibility presented on a log scale. A dotted line defines the interval from 165–140 cm where some zero values were measured. **(c)** Organic-matter content. Black bars indicate the location and age of chronologic control points (Table 1). The dashed line on the PC1 plot shows where we have interpreted the absence of ice from the catchment during the Holocene Thermal Maximum (HTM), yellow shaded region. Blue shading defines the Neoglacial period when ice was reformed during the late Holocene (4.1 ka–present).

**Glacier response to North Atlantic climate variability during the Holocene**

N. L. Balasacio et al.

[Title Page](#)

[Abstract](#) | [Introduction](#)

[Conclusions](#) | [References](#)

[Tables](#) | [Figures](#)

[◀](#) | [▶](#)

[◀](#) | [▶](#)


[Back](#) | [Close](#)

[Full Screen / Esc](#)


[Printer-friendly Version](#)

[Interactive Discussion](#)





**Figure 4.** Regional response of glaciers to Holocene climate changes. **(a)** Kulusuk glaciers interpreted from PC1 data with July insolation anomalies at 65° N (Berger and Loutre, 1991). **(b)** Hematite-stained grains (HSG) identified in core MC52-VM29-191 interpreted to indicate ice-rafting events (Bond et al., 1997). **(c)** Ratio of total organic carbon to total nitrogen (C/N) and **(d)** changes in sedimentation rate from Hvítárvatn, interpreted to reflect changes in the size of the Langjökull ice cap, Iceland and catchment instability in response to climate cooling (Larsen et al., 2012). Yellow shading marks the timing of the regional Holocene Thermal Maximum (HTM). Blue bars highlight intervals of glacier advance and increased ice rafting that define Neoglacial cooling events.



**Figure 5.** Change in the size of the Kulusuk glaciers since AD 700 compared with other high resolution glacier and ice caps records from the region. **(a)** The Kulusuk PC1 record. Black horizontal line shows average value over this period. **(b)** Big Round Lake, Baffin Island, varve thickness and magnetic susceptibility (Thomas and Briner, 2009). **(c)** Baffin Island ice cap activity reconstructed using vegetation kill dates with text showing original interpretations (Miller et al., 2012). **(d)** Langjökull ice cap, Iceland based on varve thickness from Lake Hvítárvatn (Larsen et al., 2011). Blue shading marks periods of increased glacier size (sustained above average PC1 values).

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Glacier response to North Atlantic climate variability during the Holocene

N. L. Balasacio et al.



**Figure 6.** Relative rates of change in the size of the Kulusuk glaciers interpreted from scanning XRF PC1 data. Red bars show 105 year intervals when the average rate was positive indicating glacier retreat, and blue bars show intervals when the average rate was negative indicating glacier retreat. Values not calculated during the mid-Holocene when we interpret glaciers to be absent.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

